ForPatients

by Roche

Non-Small Cell Lung Cancer (NSCLC) Non Small Cell Lung Carcinoma

A Study to Evaluate Participant and Healthcare Professional Reported Preference for Subcutaneous Atezolizumab Compared With Intravenous Atezolizumab Formulation in Participants With Non-Small Cell Lung Cancer

Trial Status Trial Runs In Trial Identifier
Completed 12 Countries NCT05171777 MO43576

The source of the below information is the publicly available website ClinicalTrials.gov. It has been summarised and edited into simpler language.

Trial Summary:

This is a Phase II, randomized, multi-center, multinational, open-label, cross-over study in adult participants with PD-L1-positive NSCLC. Two populations will be included: participants with resected Stage II, IIIA, and selected IIIB (T3-N2) NSCLC who have completed adjuvant platinum-based chemotherapy without evidence of disease relapse/recurrence, and chemotherapy-naïve participants with Stage IV NSCLC. The study will evaluate participant- and healthcare professionals (HCP)-reported preference for atezolizumab subcutaneous (SC) compared with atezolizumab intravenous (IV).

Hoffmann-La Roche Sponsor		Phase 2 Phase		
NCT05171777 MO43576 Trial Identifiers				
Eligibility Criteria:				
Gender All	Age >=18 Years		Healthy Volunteers	